

Field-theory computation of charge- density oscillations in a Tomonaga-Luttinger model with impurities

Victoria I. Fernández^{a,b} and Carlos M. Naón^{a,b}

March 2000

Abstract

We compute the charge-density oscillations in a Tomonaga-Luttinger model with impurities for a short-range electron-electron potential.

Pacs: 05.30.Fk
11.10.Lm
71.10.Pm

^a Depto. de Física. Universidad Nacional de La Plata. CC 67, 1900 La Plata, Argentina.

^b CONICET.

E-Mail: victoria@venus.fisica.unlp.edu.ar

naon@venus.fisica.unlp.edu.ar

In a recent paper [1] we introduced an extended version of the Non-local Thirring model (NLT) [2] which includes a local interaction between the fermion current and a classical background field C_μ . Thus, the NLT interaction Lagrangian density

$$\mathcal{L}_{int} = \frac{1}{2}g^2 \int d^2y J_\mu(x) V_{(\mu)}(x, y) J_\mu(y), \quad (1)$$

is now modified by adding

$$\mathcal{L}_{imp} = \bar{\Psi} \not{C} \Psi - M(x) \bar{\Psi} \Psi \quad (2)$$

The name $\mathcal{L}_{imp}$ refers to the fact that this density can be used to study the interaction between electrons and impurities [3]. Indeed, if one uses (1) in order to describe the forward scattering of one-dimensional spinless electrons [2] [4], then the first (second) term in the rhs of (2) models the forward (backward) scattering between electrons and impurities. In particular, if we choose $C_1(x) = 0$ together with

$$C_0(x) = V\delta(x_0)\delta(x_1 - d) = M(x) \quad (3)$$

where V is a constant, our model coincides with the one recently considered in ref.[5] to study Friedel charge-density oscillations in a 1d Tomonaga-Luttinger liquid [6] with an impurity located at $x_1 = d$.

We now start by considering the general case ($C_\mu(x)$ and $M(x)$ arbitrary). Using a convenient representation of the functional delta and introducing the vector field A_μ (Please see ref. [7] for details), the partition function of the model under consideration can be written as

$$Z = \int DA_\mu \det(i\not{\partial} + i\gamma_0 p_F + g\not{A} + \not{C} - M(x)) e^{-S[A]}, \quad (4)$$

with

$$S(A) = \frac{1}{2} \int d^2x d^2y V_{(\mu)}^{-1}(x, y) A_\mu(x) A_\mu(y), \quad (5)$$

where $V_{(\mu)}^{-1}$ is such that

$$\int d^2z V_{(\mu)}^{-1}(z, x) V_{(\mu)}(y, z) = \delta^2(x - y). \quad (6)$$

As shown in [1] the partition function (4) is equivalent to a purely bosonic one, corresponding to a non-local Sine-Gordon model given by

$$\mathcal{L}_{NLSG} = \frac{1}{2}(\partial_\mu \varphi(x))^2 + \frac{1}{2} \int d^2 y \partial_\mu \varphi(x) d_{(\mu)}(x, y) \partial_\mu \varphi(y) + \mathcal{L}'_{imp}, \quad (7)$$

with

$$\mathcal{L}'_{imp} = F_\mu(x) \partial_\mu \varphi(x) - \frac{\alpha_0(x)}{\beta^2} \cos \beta \varphi(x) \quad (8)$$

$F_\mu(x)$ represents a couple of classical functions to be related to the C_μ 's and $\alpha_0(x)$ is to be related, of course, to $M(x)$. Indeed, it is straightforward to show that the corresponding partition functions coincide provided that the following identities hold:

$$\frac{1}{\frac{g^2}{\pi}(p_0^2 \hat{V}_{(1)} + p_1^2 \hat{V}_{(0)}) + p^2} = \frac{\beta^2}{4\pi(p^2 + \hat{d}_{(0)}p_0^2 + \hat{d}_{(1)}p_1^2)}, \quad (9)$$

$$M(x) \cos(2p_F x_1) = \frac{\alpha_0(x)}{\beta^2} \quad (10)$$

and

$$\begin{aligned} 2i\hat{V}_{(\mu)}^{-1}(-p)\hat{C}_\mu(-p)\frac{p_\mu C(p) - 2\epsilon_{\mu\nu}p_\nu B(p)}{\Delta(p)} - \\ \frac{2g^2 p_\mu p_\mu B(p)\hat{h}(-p)}{\Delta(p)} = \\ -\beta \frac{\hat{F}_\mu(-p)p_\mu}{p^2 + \hat{d}_{(0)}p_0^2 + \hat{d}_{(1)}p_1^2} \end{aligned} \quad (11)$$

where

$$A(p) = \frac{g^2}{2\pi} p^2 + \frac{1}{2}[\hat{V}_{(0)}^{-1}(p)p_1^2 + \hat{V}_{(1)}^{-1}(p)p_0^2], \quad (12)$$

$$B(p) = \frac{1}{2}[\hat{V}_{(0)}^{-1}(p)p_0^2 + \hat{V}_{(1)}^{-1}(p)p_1^2], \quad (13)$$

$$C(p) = [\hat{V}_{(0)}^{-1}(p) - \hat{V}_{(1)}^{-1}(p)]p_0p_1, \quad (14)$$

$$\Delta(p) = C^2(p) - 4A(p)B(p), \quad (15)$$

and

$$D(p, x_i, y_i) = \sum_i (e^{ipx_i} - e^{ipy_i}). \quad (16)$$

As stated in the introductory paragraph, the path-integral identification depicted above allows to make contact with recent descriptions of 1d electronic systems in the presence of fixed (not randomly distributed) impurities [5]. We shall now discuss this possibility. To be specific we shall consider the charge-density expectation value in a Tomonaga-Luttinger electronic liquid ($\hat{V}_{(1)} = 0$), with a non-magnetic impurity located at $x_1 = d$,

$$\rho(x) = \langle \frac{i}{\sqrt{\pi}} \partial_1 \varphi - \cos(2p_F x_1) \cos(2\sqrt{\pi} \varphi) \rangle_{\mathcal{L}_{Bos}} \quad (17)$$

where

$$\begin{aligned} \mathcal{L}_{Bos} = & \frac{1}{2}(\partial_\mu \varphi(x))^2 + \int d^2 y \partial_1 \varphi(x) \frac{V_{(0)}(x, y)}{\pi} \partial_1 \varphi(y) \\ & - 2i \int d^2 y p_F x_1 (V_{(0)}(x, y) \partial_1^y) \partial_1 \varphi(y) \\ & + \frac{i}{\sqrt{\pi}} C_0(x) \partial_1 \varphi \\ & - C_0(x) \cos(2p_F x_1) \cos(2\sqrt{\pi} \varphi). \end{aligned}$$

It is convenient to have an even Lagrangian density. This is easily achieved through a translation in the field φ . The result is

$$\begin{aligned} \mathcal{L}'_{Bos} = & \frac{1}{2}(\partial_\mu \varphi(x))^2 \\ & + \int d^2 y \partial_1 \varphi(x) \frac{V_{(0)}(x, y)}{\pi} \partial_1 \varphi(y) \\ & - C_0(x) \cos(2p_F x_1) \cos(2\sqrt{\pi} \varphi) \end{aligned} \quad (18)$$

where f is a classical function satisfying

$$\partial_0 f = 0 \quad (19)$$

and

$$\begin{aligned} & \partial_1 f + \frac{2}{\pi} \int d^2 y V_{(0)}(x, y) \partial_1^y f \\ & + \frac{i}{\sqrt{\pi}} C_0 + 2ip_F \int d^2 y y_1 \partial_1^x V_0(x, y) = 0 \end{aligned} \quad (20)$$

Thus we get

$$\begin{aligned} \rho(x) - \frac{i}{\sqrt{\pi}} \partial_1 f &= -\cos(2p_F x_1) \cos(\sqrt{4\pi} f(x_1)) \\ & \langle \cos(\sqrt{4\pi} \varphi) \rangle_{\mathcal{L}'_{Bos}}, \end{aligned} \quad (21)$$

where $\cos(2p_F x_1) \cos(\sqrt{4\pi} f(x_1))$ is called the Friedel oscillation and $A(x) = \langle \cos(\sqrt{4\pi} \varphi) \rangle_{\mathcal{L}'_{Bos}}$ is the corresponding envelope.

Since $\mathcal{L}'_{Bos}$ is not exactly solvable, we shall use the well-known self consistent harmonic approximation (SCHA) [12], which consists in replacing $\mathcal{L}'_{Bos}$ by

$$\begin{aligned} \mathcal{L} &= \frac{1}{2} (\partial_\mu \varphi(x))^2 + \frac{m(x)}{2} \varphi^2 + \\ & \frac{1}{\pi} \int dy_1 \partial_1^x \varphi(x_0, x_1) U(x_1 - y_1) \partial_1^y \varphi(x_0, y_1) \end{aligned} \quad (22)$$

where $m(x)$ is an arbitrary function, related to the impurity, to be variationally determined. Also note that we have already considered the most interesting case of instantaneous potentials of the form $V_{(0)}(x, y) = \delta(x_0 - y_0) U(x_1 - y_1)$.

Let us now compute the envelope $A(x)$ in this approximation:

$$A(x) = \langle e^{-\int d^2 y J(x, y) \varphi(y)} \rangle_{\mathcal{L}} \quad (23)$$

with

$$J(x, y) = -2i\sqrt{\pi} \delta^2(x - y) \quad (24)$$

We are interested in the case

$$m(x) = m\delta(x_1 - d). \quad (25)$$

Going to momentum space and doing a translation in the field $\hat{\varphi}(p) \rightarrow \hat{\phi}(p) + a(p)$, with $a(p)$ a classical function, we find

$$A(x) = \exp -\frac{1}{2} \int \frac{d^2 p}{(2\pi)^2} J(-p) a(p) = e^{i\sqrt{\pi}a(x)} \quad (26)$$

where $a(p)$ satisfies

$$a(p) = \frac{-J(p)}{2R(-p)} - \int \frac{d^2 q}{(2\pi)^2} m(p-q) \frac{a(q)}{2R(-p)} \quad (27)$$

with

$$\begin{aligned} R(p) &= \frac{p^2}{2} + p_1^2 \frac{U(p_1)}{\pi} \\ m(k) &= 2\pi m \delta(-k_0) e^{-ik_1 d} \\ J(k) &= -2i\sqrt{\pi} e^{-ik \cdot x} \end{aligned} \quad (28)$$

As we can see, the calculation of $A(x)$ is reduced to finding the classical function $a(p)$, that is, to solving the integral equation (27). This is an integral equation with degenerated kernel whose solution is [13]

$$a(p) = \frac{-J(p)}{2R(p)} + K(p_0) f(-p) \quad (29)$$

with

$$\begin{aligned} f(p) &= -\frac{m}{2} \frac{e^{ip_1 d}}{R(p)} \\ g(q_1) &= \frac{e^{iq_1 d}}{2\pi} \\ K(-p_0) &= \frac{-\frac{1}{2} \int_{-\infty}^{\infty} dq_1 g(q_1) \frac{J(-p_0, q_1)}{R(p_0, q_1)}}{1 - \int_{-\infty}^{\infty} dq_1 g(q_1) f(p_0, -q_1)} \end{aligned} \quad (30)$$

It is straightforward to verify that $a(p)$ can be written in a more compact way as

$$a(p) = \frac{i\sqrt{\pi}}{R(p)} e^{-ip \cdot x} \left[1 - \frac{me^{ip_1 r} I(p_0, r)}{\pi + mI(p_0, 0)} \right] \quad (31)$$

where

$$I(p_0, r) = \int_0^\infty dq_1 \frac{\cos(q_1 r)}{p_0^2 + q_1^2 (1 + \frac{2U(q_1)}{\pi})}. \quad (32)$$

with $r = |d - x_1|$.

So we finally obtained $A(r)$ as a functional of the electron-electron potential:

$$A(r) = \exp -\frac{1}{\pi} \int_{-\infty}^\infty dp_0 \left(I(p_0, 0) - \frac{mI^2(p_0, r)}{\pi + mI(p_0, 0)} \right). \quad (33)$$

This is the main formal result of our work. In order to check the validity of our formula we shall now consider a short-range interaction $U(q_1) = U = \text{constant}$. It is convenient to express $A(r)$ in the form

$$A(r) = \exp(T + W)(r) \quad (34)$$

where one has defined

$$\begin{aligned} T &= -\frac{2}{\pi} \int_0^\infty dp_0 I(p_0, 0) \\ W(r) &= \frac{m}{\pi} \int_{-\infty}^\infty dp_0 \frac{I^2(p_0, r)}{\pi + mI(p_0, 0)} \end{aligned} \quad (35)$$

These integrals are easily performed yielding

$$\begin{aligned} T &= -\frac{2}{\pi} \int_0^\infty dp_0 \int_0^\infty dq_1 \frac{1}{p_0^2 + q_1^2 (1 + \frac{2U}{\pi})} \\ &= \ln\left[\frac{\mu}{\Lambda}\right]^\lambda \end{aligned} \quad (36)$$

and

$$\begin{aligned}
W &= \frac{m}{\pi} \int_{-\infty}^{\infty} dp_0 \frac{I^2(p_0, r)}{\pi + mI(p_0, 0)} \\
&= -\frac{1}{\lambda} e^{\frac{2\mu r}{\lambda}} E_i\left(-\frac{2\mu r}{\lambda}\right) \\
&\quad + \frac{1}{\lambda} e^{\frac{mr}{\lambda^2}} E_i\left(-\frac{mr}{\lambda^2}\right),
\end{aligned} \tag{37}$$

where $E_i(x)$ is the exponential integral function, $\mu \rightarrow 0^+$ and $\Lambda \rightarrow \infty$ are infrared and ultraviolet cutoffs respectively, and $\lambda = \sqrt{1 + \frac{2U}{\pi}}$.

At this point, in order to compare our results with previous computations we consider the long distance regime defined by $\frac{2\mu r}{\lambda} \ll 1$, $\frac{mr}{\lambda^2} \gg 1$. We get

$$A(r) = C(\lambda, \mu, \Lambda) \frac{\lambda^{\frac{1}{\lambda}}}{2r} e^{-\frac{\lambda}{mr}} \tag{38}$$

which exactly coincides with [11] and [5] under the same regime.

Acknowledgements

The authors are supported by Universidad Nacional de La Plata (UNLP) and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina. They are grateful to the organizers of the XX Encontro Nacional de Física de Partículas e Campos (Brazil) where these results were presented.

References

- [1] V. Fernández, K. Li, C. Naón, Physics Letters B **452**, 98, (1999).
- [2] C.Naón, M.C.von.Reichenbach, M.L. Trobo, Nucl. Phys. **B 435** [fs], 567, (1995).
- [3] C.L.Kane, M.P.A.Fisher, Phys.Rev.B **46** 15, 233, (1992).
- [4] J.Voit, Rep. on Progr. in Phys. **58**, 977, (1995).
- [5] Q. Yuan, H.Chen, Y.Zhang, Y.Chen, Phys.Rev.B **58**, 1084, (1998).
- [6] S.Tomonaga, Prog. Theor. Phys. **5**, 544, (1950).
J.Luttinger, J. Math. Phys. **4**, 1154, (1963).
E.Lieb and D.Mattis, J. Math. Phys. **6**, 304, (1965).
- [7] Kang Li, Carlos Naón, J. Phys. A **31**, 7929, (1998).
- [8] H.J.Schulz, Phys.Rev.Lett. **71**, 1864, (1993).
- [9] K.Fujikawa, Phys. Rev. D **21**, 2848, (1980).
- [10] M.Manías, C.Naón, M.L.Trobo, Physics Letters B **416**, 157, (1998).
- [11] R. Egger, H. Grabert, Phys.Rev.Lett. **75**, 3505, (1995).
- [12] Y. Saito, Z. Phys. B **32**, 75, (1978).
M. Stevenson, Phys. Rev. D **32**, 1389, (1985).
M.P.A. Fisher, W. Zwerger, Phys. Rev. B **32**, 6190, (1985).
A.O. Gogolin, Phys. Rev. Lett. **71**, 2995, (1993).
N.V. Prokof'ev, Phys. Rev. B **49**, 2243, (1994).
B.W. Xu, Y.M. Zhang, J. Phys. A **29**, 7349, (1996).
- [13] See for instance: S.L. Sobolev, "Partial Differential Equations of Mathematical Physics", Dover Pub. Inc., 1989.